


$$x_d(t) = x(t) + \frac{d y'(t)}{\sqrt{x'(t)^2 + y'(t)^2}}$$


$$y_d(t) = y(t) - \frac{d x'(t)}{\sqrt{x'(t)^2 + y'(t)^2}}$$



Расстояние d от точки $M_0(x_0, y_0)$ до прямой, заданной уравнением общего вида $Ax + By + C = 0$ определяется по формуле:



$$d = \frac{|Ax_0 + By_0 + C|}{\sqrt{A^2 + B^2}}$$

общее
 $Ax + By + C = 0$
 $Ax = -(By + C)$
каноническое
 $Ax / (-B) = (y + C/B)$
 $x / (-B) = (y + C/B) / A = t$
параметрическое
 $x = -Bt$
 $y = At + C/B$




$L = \int V(dx^2 + dy^2)$
 $y = y(t) \quad dy/dt = y'(t) \quad dy = y'(t)dt$
 $x = x(t) \quad dx/dt = x'(t) \quad dx = x'(t)dt$

$S[a; b] \int V(dx^2 + dy^2) = S[a; b] \int V(1 + (dy/dx)^2) dx$
 $S[a; b] \int V(dx^2 + dy^2) = S[a; b] \int V(y'(t)^2 + x'(t)^2) dt$




$n\{y'(t), -x'(t)\} \cdot g = s\{w - x(t), q - y(t)\}$
 $w = x(t) + g \cdot y'(t)$
 $q = y(t) - g \cdot x'(t)$



$a^2 = (w - x(t))^2 + (q - y(t))^2$
 $[w - x(t)] / y'(t) = [q - y(t)] / -x'(t)$

$n\{y'(t), -x'(t)\}$
 $|n| = \sqrt{[y'(t)]^2 + [-x'(t)]^2} = \sqrt{y'(t)^2 + x'(t)^2}$
 $e = n/|n| = \{y'(t)/\sqrt{y'(t)^2 + x'(t)^2}, -x'(t)/\sqrt{y'(t)^2 + x'(t)^2}\}$
 $E\{a \cdot y'(t)/\sqrt{y'(t)^2 + x'(t)^2}, -a \cdot x'(t)/\sqrt{y'(t)^2 + x'(t)^2}\}$



$w - x(t) = a \cdot y'(t) / \sqrt{y'(t)^2 + x'(t)^2}$
 $q - y(t) = -a \cdot x'(t) / \sqrt{y'(t)^2 + x'(t)^2}$
 $w = x(t) + a \cdot y'(t) / \sqrt{y'(t)^2 + x'(t)^2}$
 $q = y(t) - a \cdot x'(t) / \sqrt{y'(t)^2 + x'(t)^2}$