

любое тригур-ие можно свести к ур-ию высших степеней

$$\sin^2 x = \sin^2 x / 1 = \sin^2 x / (\sin^2 x + \cos^2 x) = \tan^2 x / (1 + \tan^2 x)$$

$$\cos^2 x = \cos^2 x / (\cos^2 x + \sin^2 x) = 1 / (1 + \tan^2 x)$$

$$\begin{aligned} \sin 2x &= \sin 2x / 1 = \sin 2x / (\cos^2 x + \sin^2 x) = 2 \sin x \cos x / (\cos^2 x + \sin^2 x) = \\ &= 2 \tan x / (1 + \tan^2 x) \end{aligned}$$

$$\cos 2x = \cos 2x / 1 = (\cos^2 x - \sin^2 x) / (\sin^2 x + \cos^2 x) = (1 - \tan^2 x) / (1 + \tan^2 x)$$

$$\begin{aligned} \tan 2x &= \sin 2x / \cos 2x = 2 \tan x / (1 + \tan^2 x) / (1 - \tan^2 x) / (1 + \tan^2 x) = \\ &= 2 \tan x / (1 - \tan^2 x) \end{aligned}$$

$$\tan 3x = \tan(x + 2x) = (\tan x + \tan 2x) / (1 - \tan x \tan 2x)$$

$$\begin{aligned} \tan(a+b) &= \sin(a+b) / \cos(a+b) = (\sin a \cos b + \sin b \cos a) / (\cos a \cos b - \sin a \sin b) \\ &= (\tan a + \tan b) / (1 - \tan a \tan b) \end{aligned}$$

проверить отдельно

$$\cos x = 0$$

$$3 \tan 3x - \cot 2x = 4 \tan x$$